

Macroinvertebrates & Land Use Change: Pollution Tolerance

Time: 2 class periods

National Benchmarks: Benchmarks 5A: Diversity of Life; 5D Interdependence of Life; 5E: Flow of Matter and Energy; 9B:Symbolic Relationships; 9D:Uncertainty; 12B:Computation and Estimation; 12D:Communication Skills; 12E:Critical-Response Skills.

National Science Content Standards: *Science as Inquiry: A; Life Science: C:* Biological Evolution; The Interdependence of Organisms; Matter, Energy, and Organization in Living Systems; *Science and Technology: E:* Abilities of Technological Design; Understandings about Science and Technology; *Science in Personal and Social Perspectives: F:* Population Growth; Natural Resources: Environmental Quality; Natural and Human-induced Hazards; Science and Technology in Local, National, and Global Challenges

New York State Standards: 1, 2, 3, 4, 5, 6, 7

Objective: Students will know how land use affects water quality, and be able to use macroinvertebrates to understand the impact of land use change in watersheds.

Lesson Outline:

1. Students select watersheds for the study, and create hypotheses for macroinvertebrates based on observed land use.
2. Students obtain aerial photos of the study site, and determine amount of impervious surface in each tested watershed.
3. Students complete macroinvertebrate studies at two different sites, looking at the pollution tolerance values of the invertebrates, and compile and discuss results.
4. Students compare their results with published scientific data.

Materials

Historic and current aerial photos (from website)

Aerial photos or Google maps of watershed(s)

Transparency with grid

Waders or appropriate shoes

Leaf pack bags (plastic mesh bags for leaf litter)

Dissecting trays, tweezers, sorting containers to observe benthic material

Test kits for DO, phosphates, nitrates, pH, chloride and other appropriate tests (optional)

Goggles, gloves

Data sheets for macroinvertebrates

Preparation: Aerial photos are a great way for students to compare land use types. Local extension offices or a university GIS department may have maps you can use, although you can also print aerial photos directly from a web application like Google Maps. Depending on the level of your students, you may want to identify the test watersheds ahead of time. Determining watershed boundaries is easy to do using contour lines, which are shown in the terrain feature within Google Maps (the online version, not Google Earth, which does not have contour lines). Google Earth has a tremendous 3D view of terrain, which students can also use to delineate watershed boundaries. Once the watersheds are identified, print them out for students to calculate the different types of land use.

Engage: Ask students to identify the major land use type in their neighborhood. How does land use change when they drive in different directions? Is land use the same everywhere?

Provide students with the historic and current aerial photos, and ask them to identify the major trends evident in the photos. You can also use the accompanying powerpoint to show pictures of land use from other parts of the country and the world.

Explore: Students should begin working on their experimental set-up.

- 1) Students identify test watersheds A & B (unless you do this beforehand).
- 2) Students determine % land cover of their test watersheds.
- 3) Students make a prediction, using a provided graph, about the results of a watershed comparison study.
- 4) Students collect macroinvertebrates. There are a variety of methods for collecting these organisms, which can be found in the accompanying document produced by Hudson Basin River Watch. Alternatively, you can set out leaf packs in the different watersheds and collect them back in 2-3 weeks. For leaf pack methods, visit the Stroud Center's Leaf Pack website: www.stroudcenter.org/lpn/index.htm. The Leaf Pack Network has a large range of resources available for use.
- 5) Students compile their data and identify the numbers of groups that they found in the two different watersheds. Students calculate "species" evenness and richness, although what they're calculating is actually group evenness and richness, since this lesson does not identify to species level.
- 6) Students calculate the pollution tolerance levels of each watershed, using the accompanying data sheet.
- 7) Students graph class data. If you have different watersheds for each student group, you will see a better trend than if the groups all did the same watersheds.
- 8) Students compare their data with data from other scientific sources.

Explain: After you return to the classroom, discuss student findings. What did students notice? Ask students to think about the connections between the organisms that live in/near the aquatic ecosystem with the land use in the ecosystem's watershed.

Table 1: Comparison of One Acre of Parking Lot Versus One Acre of Meadow in Good Condition

Runoff or Water Quality Parameter	Parking Lot	Meadow
Curve number (CN)	98	58
Runoff coefficient	0.95	0.06
Time of concentration (minutes)	4.8	14.4
Peak discharge rate (cfs), 2 yr., 24 hr. storm	4.3	0.4
Peak discharge rate (cfs), 100 yr. storm	12.6	3.1
Runoff volume from one-inch storm (cubic feet)	3450	218
Runoff velocity @ 2 yr. storm (feet/second)	8	1.8
Annual phosphorus load (lbs/ac./yr.)	2	0.50
Annual nitrogen load (lbs/ac./yr.)	15.4	2.0
Annual zinc load (lbs/ac./yr.)	0.30	ND


Key Assumptions:

Parking lot is 100% impervious with 3% slope, 200 feet flow length, Type 2 Storm, 2 yr. 24 hr. storm = 3.1 inches, 100 yr. storm = 8.9 inches, hydraulic radius = 0.3, concrete channel, and suburban Washington 'C' values.

Meadow is 1% impervious with 3% slope, 200 foot flow length, good vegetative condition, B soils, and earthen channel.

Urbanization can be a problem for various reasons, since impervious surfaces change the volume and the timing of runoff, and there may be contaminants in the water. In urban areas surface runoff carries pollutants from substances that have leaked or spilled onto the ground, such as oil or salt. Other pollutants such as nitrogen or phosphorus also accumulate in waterways. If you are interested in this topic, have students complete the "Land Use and Water Quality" lesson, which includes data on the increasing amount of nitrogen in suburban waterways. In addition to contamination, the amount of runoff increases with urbanization. In many cities and towns, rainwater goes into a sewage treatment plant along with sewage. When there is a heavy storm, however, and the sewage system becomes

overwhelmed, it cannot treat the water, and it often releases *all* the water into the nearest water body. This is called “combined sewage overflow” (CSO). For more information on CSOs, use the “History of Wastewater” reading.



From Schueler & Holland.

Streams and rivers across the country have been artificially channeled, straightened, or otherwise altered. Construction or poor land planning causes excess sediment to wash into streams and rivers, causing them to fill up prematurely, which adds to the flood threat. Streams normally flood, so trying to stop them from flooding in one place can cause ecological damage as well as increase flood damage in another. Urbanized streams that receive a large amount of water in a short amount of time during a storm are called “flashy” streams.

Macroinvertebrates are an important indicator of the health of an aquatic ecosystem. Immature insects such as stoneflies, mayflies, and water pennies (a type of beetle larvae) require a high amount of dissolved oxygen (DO), while aquatic worms, leeches and pond snails can survive in water with low DO. Oxygen-loving species like mayflies and stoneflies “indicator species” because they provide important clues about the water they are living in. If you only find animals like leeches, snails, and aquatic worms, then you know that there is a problem with water quality, and you should do additional studies to determine the cause.

One of the most useful indicators is the diversity or numbers of kinds of organisms. If you find only one or two kinds of animals, no matter what kind they are, you should perform other water quality tests to determine what might be wrong with your aquatic ecosystem. Sometimes, low diversity can indicate a pollution problem or other habitat change that is affecting the ecosystem.

A number of factors besides imperviousness can influence the diversity and density of macroinvertebrates present in an aquatic ecosystem. Seasons, life cycles, types of substrate, food sources, water velocity, and sampling techniques can all affect the diversity in your sample. For example, if you are testing the water in the spring, you might find fewer animals after a flood or heavy rain.

It is also important to know the animals’ life cycles. Many larvae emerge as adults in late spring and are present only as eggs during other parts of the year. The substrate on the river bottom can affect your results as well. A rocky bottom provides more habitat than a silty or

muddy bottom. You should also take into consideration the surrounding habitat: a forest often provides more food (in the form of plant material) than a meadow. Finally, you need to decide what kind of sampling technique you are going to use. A screen or net that is too large will cause you to miss some animals, while inappropriate equipment use means you won't collect a good sample of all the animals living in the ecosystem.

Extend: Students can create a presentation of their research for community members or another audience within the school, and discuss ways of improving water quality through land use change or specific mitigation strategies (pervious asphalt, rain gardens, riparian zones, etc).

Evaluate: Students turn in the lab report.

Background information:

Table 3: A Possible Scheme for Classifying and Managing for Headwater Urban Streams Based on Ultimate Imperviousness			
Urban Stream Classification	Sensitive (0-10% Imperv.)	Impacted (11-25% Imperv.)	Non-supporting (26-100% Imperv.)
Channel stability	Stable	Unstable	Highly Unstable
Water quality	Good	Fair	Fair-Poor
Stream biodiversity	Good-Excellent	Fair-Good	Poor
Resource objective	Protect biodiversity and channel stability	Maintain critical elements of stream quality	Minimize downstream pollutant loads
Water quality objectives	Sediment and temperature	Nutrient and metal loads	Control bacteria
Stormwater Practice Selection Factors	Secondary environmental impacts	Removal efficiency	Removal efficiency
Land Use Controls	Watershed-wide imp. cover limits (ICLs), site ICLs	Site imp. cover limits (ICLs)	Additional infill and redevelopment encouraged
Monitoring and enforcement	GIS monitoring of imp. cover, biomonitoring	Same as "Stressed"	Pollutant load modeling
Development rights	Transferred out	None	Transferred in
Riparian buffers	Widest buffer network	Average bufferwidth	Greenways

From: Schueler & Holland




Figure from: www.stormwatercenter.net

The model classifies streams into one of three categories: sensitive, impacted, and non-supporting. Each stream category can be expected to have unique characteristics as follows:

Sensitive Streams. These streams typically have a watershed impervious cover of zero to 10 percent. Consequently, sensitive streams are of high quality, and are typified by stable channels, excellent habitat structure, good to excellent water quality, and diverse communities of both fish and aquatic insects. Since impervious cover is so low, they do not experience frequent flooding and other hydrological changes that accompany urbanization. It should be noted that some sensitive streams located in rural areas may have been impacted by prior poor grazing and cropping practices that may have severely altered the riparian zone, and consequently, may not have all the properties of a sensitive stream. Once riparian management improves, however these streams are often expected to recover.

Impacted Streams. Streams in this category possess a watershed impervious cover ranging from 11 to 25 percent, and show clear signs of degradation due to watershed urbanization. The elevated storm flows begin to alter stream geometry. Both erosion and channel widening are clearly evident. Stream banks become unstable, and physical habitat in the stream declines noticeably. Stream water quality shifts into the fair/good category during both storms and dry weather periods. Stream biodiversity declines to fair levels, with most sensitive fish and aquatic insects disappearing from the stream.

Non-Supporting Streams. Once watershed impervious cover exceeds 25%, stream quality crosses a second threshold. Streams in this category essentially become conduits for conveying stormwater flows, and can no longer support a diverse stream community. The stream channel becomes highly unstable, and many stream reaches experience severe widening, downcutting, and streambank erosion. Pool and riffle structure needed to sustain fish is diminished or eliminated and the substrate can no longer provide habitat for aquatic insects, or spawning areas for fish. Water quality is consistently rated as fair to poor, and water recreation is no longer possible due to the presence of high bacterial levels. Subwatersheds in the non-supporting category will generally display increases in nutrient loads to downstream receiving waters, even if effective urban BMPs are installed and maintained. The biological quality of non-supporting streams is generally considered poor, and is dominated by pollution tolerant insects and fish.

References:

- Behar, S. and M. Cheo. 2004. *Hudson Basin River Watch Guidance Document*. River Network. www.hudsonbasin.org
- Environmental Indicators Worksheets*. Center for Watershed Protection. Retrieved 4/22/2008 at www.stormwatercenter.net
- Groffman, P., Law, N., Belt, K., Band, L., and G. Fisher. 2004. Nitrogen Fluxes and Retention in Urban Watershed Ecosystems. *Ecosystems* 7:393-403.
- Limburg, K.E. & R.E. Schmidt. 1990. Patterns of Fish Spawning in Hudson River Tributaries: Response to an Urban Gradient? *Ecology*, 71(4): 1238-1245.
- Morgan, R.P. & S.F. Cushman. 2005. Urbanization effects on stream fish assemblages in Maryland. *Journal of North American Benthological Society*, 24(3):643-655
- Schueler, T. R. & H.K. Holland, eds. 2000. The Importance of Imperviousness. *Watershed Protection Techniques*, 1(3): 100-111.
- Wang, L. and P. Kanehl. 2003. Influences of Watershed Urbanization and Instream Habitat on Macroinvertebrates in Cold Water Streams. *Journal of the American Water Resources Association*, 39(5): 1181-1196.